




BioDea

## CULTIVATION SHEET

*"it includes about 13000 species, including mainly herbaceous plants, but also trees, shrubs or lianas"*

Family: Compositae



some examples: wormwood,  
artichoke, chamomile, thistle,  
chicory, tarragon

SOWING TREATMENT fertigation	DOSE
IN SEEDBED: Start by wetting the substrate when sowing and continue until the moment of transplantation, with interventions every 4 days	1:800 or 125 ml/hl
TRANSPLANTATION TREATMENT fertigation	DOSE
During the transplantation of the plant 4th-6th leaf	1:500 or 200 ml/hl
FOLIAR TREATMENTS	DOSE
4th/6th leaf	1:600 or 167 ml/hl
After the 7th leaf	1:500 or 200 ml/hl
During the vegetative development until harvesting	1:500 or 200 ml/hl
<i>Repeat every 6/7 days based on seasonal conditions and to make the environment unwelcome to insects and parasites in the most critical periods, perform the foliar treatment after each mowing.</i>	
FERTIGATION	DOSE
4th/6th leaf	1:500 or 200 ml/hl
After the 7th leaf	1:300 or 333 ml/hl repeat every 14 days
<i>Repeat every 10-15 days based on needs distribute in the last irrigation phase</i>	

**Compatibility:**

The product has no contraindications for miscibility with other formulations. It is advisable to carry out compatibility tests on crops on small surfaces before extending the treatment to larger areas.

**Warnings:**

To be used exclusively in agriculture. Whoever uses the product is responsible for any damage resulting from the improper use of the product. Compliance with the aforementioned instructions is an essential condition to ensure the effectiveness of the treatment and to avoid damage to plants, people, animals.