



BioDea

CULTIVATION SHEET

"key feature is the arrangement of flowers in an umbrella shape, hence their name. Some species, such as hemlock, can be toxic, while others are edible"

Family: Umbelliferae



some examples: *dill, anise, angelica, carrot, caraway caraway, celery, chervil, coriander, caraway, fennel, parsnip, parsley*

TREATMENT DURING SOWING IN FERTIGATION	DOSE
Start by wetting the soil when sowing, with interventions every 3-4 days	1:800 or 125 ml/hl
FOLIAR TREATMENT	DOSE
5th/6th leaf	1:600 or 167 ml/hl
After the 6th leaf	1:500 or 200 ml/hl
From the growth of the tuber or of the root until harvesting	1:500 or 200 ml/hl
<i>Repeat every 6/7 days based on seasonal conditions and to make the environment unwelcome to insects and parasites in the most critical periods</i>	
FERTIGATION	DOSE
5th/6th leaf	1:500 or 200 ml/hl
After the 6th leaf until the harvest of the tuber or of the root	1:300 or 333 ml/hl repeat every 14 days
<i>Repeat every 10-15 days based on needs distribute in the last irrigation stage</i>	

Compatibility:

The product has no contraindications for miscibility with other formulations.
It is advisable to carry out compatibility tests on crops on small surfaces before extending the treatment to larger areas.

Warnings:

To be used exclusively in agriculture.
Whoever uses the product is responsible for any damage resulting from the improper use of the product.
Compliance with the aforementioned instructions is an essential condition to ensure the effectiveness of the treatment and to avoid damage to plants, people, animals.